Page 2 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

## Amendments to the Claims

This listing of claims replaces all prior versions, and listings, of claims in the above-identified application:

- 1. (Previously Presented) A water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkyl-Y-containing side chains, wherein Y is O or NR, wherein R is H or CH<sub>3</sub>, and wherein the alkyl group of the alkyl-Y-containing side chain has at least 4 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- (Original) The water-in-oil emulsion of claim 1 wherein the vinyl polymer is soluble in the oil phase.
- (Original) The water-in-oil emulsion of claim 1 wherein the ethylene oxide groups and alkyl-Y groups are in different side chains.
- 4. (Original) The water-in-oil emulsion of claim 1 which is stable.
- 5. (Original) The water-in-oil emulsion of claim 1 which is substantive.
- 6. (Previously Presented) The water-in-oil emulsion of claim 5 which provides a reduction in skin capacitance of greater than about 15% compared to an untreated portion of the skin.

Page 3 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

- 7. (Original) The water-in-oil emulsion of claim 1 wherein the ethylene oxidecontaining side chains further include isopropylene oxide groups.
- 8. (Original) The water-in-oil emulsion of claim 1 wherein the ethylene oxidecontaining side chains include at least four ethylene oxide groups.
- 9. (Original) The water-in-oil emulsion of claim 1 wherein the oil phase comprises one or more oils present in a total amount of at least about 20 wt-%, based on the total weight of the emulsion.
- 10. (Original) The water-in-oil emulsion of claim 1 wherein the ethylene oxidecontaining side chains are derived from one or more monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomers.
- 11. (Original) The water-in-oil emulsion of claim 10 wherein the monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomers have the formula:

wherein:

m is at least 2;

p is 0 to 50;

q is 0 or 1;

R<sup>3</sup> is H or CH<sub>3</sub>; and

R<sup>4</sup> is hydrogen or linear or branched alkyl and/or aryl groups; with the proviso that the isopropylene oxide groups (the "p" groups) and

Page 4 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration.

- 12. (Original) The water-in-oil emulsion of claim 1 wherein the alkyl-Y-containing side chains are derived from one or more monoethylenically unsaturated alkyl (meth)acrylic monomers.
- 13. (Original) The water-in-oil emulsion of claim 12 wherein the monoethylenically unsaturated alkyl (meth)acrylic monomers are selected from the group consisting of (meth)acrylate monomers, (meth)acrylamide monomers, and combinations thereof.
- 14. (Original) The water-in-oil emulsion of claim 12 wherein the monoethylenically unsaturated alkyl (meth)acrylic monomers are alkyl (meth)acrylate monomers having the formula:

$$\begin{array}{ccc}
R^1 & O \\
I & II \\
H_2C = C - C - OR^2
\end{array}$$

wherein:

R<sup>1</sup> is H or CH<sub>3</sub>; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms.

- 15. (Original) The water-in-oil emulsion of claim 1 further comprising a stabilizer.
- 16. (Original) The water-in-oil emulsion of claim 1 wherein the vinyl polymer is the reaction product of: about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylic monomer; and about 10 wt-% to about 40 wt-% of at least

Page 5 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer.

- 17. (Original) The water-in-oil emulsion of claim 1 which has compatibility with at least one bioactive agent.
- 18. (Original) The water-in-oil emulsion of claim 17 wherein the bioactive agent is an antimicrobial.
- 19. (Original) The water-in-oil emulsion of claim 18 wherein the antimicrobial is chlorhexidine gluconate.
- 20. (Original) The water-in-oil emulsion of claim 18 wherein the antimicrobial is an iodophor.
- 21. (Original) The water-in-oil emulsion of claim 20 wherein the iodophor is povidone-iodine.
- 22. (Original) The water-in-oil emulsion of claim 1 wherein a pressure sensitive adhesive tape applied over the emulsion on skin adheres at a level of at least about 50% of the level of adhesion of the pressure sensitive adhesive tape applied directly to the skin.
- 23. (Original) The water-in-oil emulsion of claim 1 wherein the vinyl polymer has a calculated HLB of more than about 1 and less than about 10.
- 24. (Original) The water-in-oil emulsion of claim 1 comprising at least about 0.25 wt-% of the vinyl polymer, based on the total weight of the emulsion.

Page 6 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

- 25. (Original) The water-in-oil emulsion of claim 1 comprising no more than about 10 wt-% of the vinyl polymer, based on the total weight of the emulsion.
- 26. (Original) The water-in-oil emulsion of claim 1 further comprising a humectant.
- 27. (Previously Presented) The water-in-oil emulsion of claim 1 further comprising one or more additives selected from the group consisting of humectants, surfactants, conditioners, sunscreen agents, insect repellents, vitamins, herbal extracts, antiperspirant agents, deodorant agents, skin bleaching agents, skin coloring agents, hair bleaching agents, hair coloring agents, depilating agents, antidandruff agents, antiacne agents, astringents, tensors, skin toning agents, glitter, pigments, dyes, bleaches, perfumes, fragrances, preservatives, antioxidants, waxes, film-forming polymers, propellants, buffers, organic suspending agents, inorganic suspending agents, organic thickening agents, inorganic thickening agents, plasticizers, herbal extracts, flavoring agents, com removers, callus removers, wart removers, and combinations thereof.
- 28. (Previously Presented) A water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkoxy-containing side chains, wherein the alkyl group of the alkoxy-containing side chain has 4 to 50 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 29. (Previously Presented) A water-in-oil emulsion comprising: an oil phase; a water phase; and a vinyl polymer that is insoluble or sparingly soluble in the water phase; wherein the vinyl polymer comprises the reaction product of monomers comprising:

Page 7 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylate monomer having the formula:

$$R^{1}$$
 O  
 $| | | | |$   
 $H_{7}C = C - C - OR^{2}$ 

wherein:

R1 is H or CH3; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms; and

about 10 wt-% to about 40 wt-% of at least one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer having the formula:

wherein:

m is at least 2;

p is 0 to 50;

q is 0 or 1;

R<sup>3</sup> is H or CH<sub>3</sub>; and

R<sup>4</sup> is hydrogen or linear or branched alkyl and/or aryl groups; with the proviso that the isopropylene oxide groups (the "p" groups) and the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration;

Page 8 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

with the proviso that the vinyl polymer includes no more than about 0.1 wt-% copolymerized acidic monomers.

- 30. (Original) The water-in-oil emulsion of claim 29 which is stable.
- 31. (Original) The water-in-oil emulsion of claim 29 which is substantive.
- 32. (Previously Presented) A moisturizing composition comprising a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkyl-Y-containing side chains, wherein Y is O or NR, wherein R is H or CH<sub>3</sub>, and wherein the alkyl group of the alkyl-Y-containing side chain has at least 4 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 33. (Original) The moisturizing composition of claim 32 which is stable.
- (Original) The moisturizing composition of claim 32 which is substantive.
- 35. (Original) The moisturizing composition of claim 32 wherein a pressure sensitive adhesive tape applied over the emulsion on skin adheres at a level of at least about 50% of the level of adhesion of the pressure sensitive adhesive tape applied directly to the skin.
- 36. (Previously Presented) A moisturizing composition comprising a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkoxy-containing side chains, wherein the alkyl group of the alkoxy-containing side

Page 9 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

chain has 4 to 50 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.

37. (Previously Presented) A moisturizing composition comprising a water-in-oil emulsion comprising: an oil phase; a water phase; and a vinyl polymer that is insoluble or sparingly soluble in the water phase; wherein the vinyl polymer comprises the reaction product of monomers comprising:

about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylate monomer having the formula:

$$R^{1} O \\ H_{2}C = C - C - OR^{2}$$

wherein:

R<sup>1</sup> is H or CH<sub>3</sub>; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms; and

about 10 wt-% to about 40 wt-% of at least one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer having the formula:

wherein:

Page 10 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

m is at least 2;

p is 0 to 50;

q is 0 or 1;

R<sup>3</sup> is H or CH<sub>3</sub>; and

R<sup>4</sup> is hydrogen or linear or branched alkyl and/or aryl groups; with the proviso that the isopropylene oxide groups (the "p" groups) and the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration.

- 38. (Previously Presented) A mammalian tissue antiseptic composition comprising an antimicrobial agent and a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkyl-Y-containing side chains, wherein Y is O or NR, wherein R is H or CH<sub>3</sub>, and wherein the alkyl group of the alkyl-Y-containing side chain has at least 4 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 39. (Original) The tissue antiseptic composition of claim 38 which is stable.
- 40. (Previously Presented) A mammalian tissue antiseptic composition comprising an antimicrobial agent and a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkoxy-containing side chains, wherein the alkyl group of the alkoxy-containing side chain has 4 to 50 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.

Page 11 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

41. (Previously Presented) A mammalian tissue antiseptic composition comprising: an oil phase; a water phase; an antimicrobial agent; and a vinyl polymer that is insoluble or sparingly soluble in the water phase; wherein the vinyl polymer comprises the reaction product of monomers comprising:

about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylate monomer having the formula:

$$R^1$$
 O | ||  $H_2C=C-C-OR^2$ 

wherein:

R1 is H or CH3; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms; and

about 10 wt-% to about 40 wt-% of at least one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer having the formula:

wherein:

m is at least 2;

p is 0 to 50;

q is 0 or 1;

R<sup>3</sup> is H or CH<sub>3</sub>; and

R4 is hydrogen or linear or branched alkyl and/or aryl groups;

Page 12 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

with the proviso that the isopropylene oxide groups (the "p" groups) and the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration.

- 42. (Previously Presented) A personal care composition comprising a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkyl-Y-containing side chains, wherein Y is O or NR, wherein R is H or CH<sub>3</sub>, and wherein the alkyl group of the alkyl-Y-containing side chain has at least 4 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 43. (Original) The personal care composition of claim 42 which is a hair care composition.
- 44. (Original) The personal care composition of claim 43 wherein the hair care composition is a styling agent, shampoo, dye, conditioner, rinse, antidandruff preparation, or mask for the hair.
- 45. (Original) The personal care composition of claim 42 which is in the form of an insect repellant, shaving product, hand lotion, body lotion, gel, cream, sunless tanning composition, sunscreen, cleanser, toner, astringent, freshener, mask for skin, nail polish, nail strengthener, underarm deodorant, antiperspirant, bath powder, talc, bath oil, bubble bath, makeup, cologne, perfume, composition for cushioning sores, or hair removal composition.
- 46. (Original) The personal care composition of claim 42 which is a makeup.

Page 13 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

- 47. (Original) The personal care composition of claim 46 wherein the makeup is a lipstick, eye shadow, eye liner, mascara, rouge, face powder, or foundation.
- 48. (Previously Presented) A personal care composition comprising a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkoxy-containing side chains, wherein the alkyl group of the alkoxy-containing side chain has 4 to 50 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 49. (Previously Presented) A personal care composition comprising a water-in-oil emulsion comprising: an oil phase; a water phase; and a vinyl polymer that is insoluble or sparingly soluble in the water phase; wherein the vinyl polymer comprises the reaction product of monomers comprising:

about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylate monomer having the formula:

$$R^{1}$$
 O  
 $H_{2}C = C - C - OR^{2}$ 

wherein:

R1 is H or CH3; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms; and

about 10 wt-% to about 40 wt-% of at least one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer having the formula:

Page 14 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

wherein:

m is at least 2;

p is 0 to 50;

q is 0 or 1;.

R<sup>3</sup> is H or CH<sub>3</sub>; and

R<sup>4</sup> is hydrogen or linear or branched alkyl and/or aryl groups; with the proviso that the isopropylene oxide groups (the "p" groups) and the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration.

- 50. (Previously Presented) A transdermal drug delivery composition comprising a pharmaceutical agent and a water-in-oil emulsion comprising: a vinyl polymer comprising ethylene oxide-containing side chains and alkyl Y-containing side chains, wherein Y is O or NR, wherein R is H or CH<sub>3</sub>, and wherein the alkyl group of the alkyl-Y-containing side chain has at least 4 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the vinyl polymer is insoluble or sparingly soluble in the water phase.
- 51. (Previously Presented) A transdermal drug delivery composition comprising a pharmaceutical agent and a water-in-oil emulsion comprising:

Page 15 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

a vinyl polymer comprising ethylene oxide-containing side chains and alkoxy-containing side chains, wherein the alkyl group of the alkoxy-containing side chain has 4 to 50 carbon atoms on average in a cyclic, branched-, or straight-chain configuration and optionally includes one or more heteroatoms; an oil phase; and a water phase; wherein the

vinyl polymer is insoluble or sparingly soluble in the water phase.

52. (Previously Presented) A transdermal drug delivery composition comprising a pharmaceutical agent and a water-in-oil emulsion comprising: an oil phase; a water phase; and a vinyl polymer that is insoluble or sparingly soluble in the water phase; wherein the vinyl polymer comprises the reaction product of monomers comprising:

about 60 wt-% to about 90 wt-% of at least one monoethylenically unsaturated alkyl (meth)acrylate monomer having the formula:

$$\begin{array}{c} R^1 & O \\ I & II \\ H_2C = C - C - OR^2 \end{array}$$

wherein:

R1 is H or CH3; and

R<sup>2</sup> is a linear, branched, or cyclic alkyl group optionally including one or more heteroatoms; and

about 10 wt-% to about 40 wt-% of at least one monoethylenically unsaturated poly(alkylene oxide) (meth)acrylic monomer having the formula:

$$H_2C = C - C - (OCH_2CH_2)_m(OCH_2CH)_p - O(C)_qR^4$$

Page 16 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

wherein:

m is at least 2;

p is 0 to 50;

q is 0 or 1;

R3 is H or CH3; and

R<sup>4</sup> is hydrogen or linear or branched alkyl and/or aryl groups; with the proviso that the isopropylene oxide groups (the "p" groups) and the ethylene oxide groups (the "m" groups) are arranged in a reversed, alternating, random, or block configuration.

- 53. (Withdrawn) A method of moisturizing mammalian skin comprising applying a moisturizing composition of claim 32 to mammalian skin.
- 54. (Withdrawn) A method of moisturizing mammalian skin comprising applying a moisturizing composition of claim 36 to mammalian skin.
- 55. (Withdrawn) A method of moisturizing mammalian skin comprising applying a moisturizing composition of claim 37 to mammalian skin.
- 56. (Withdrawn) A method of disinfecting mammalian tissue comprising applying a tissue antiseptic composition of claim 38 to mammalian tissue.
- 57. (Withdrawn) A method of disinfecting mammalian tissue comprising applying a tissue antiseptic composition of claim 40 to mammalian tissue.
- 58. (Withdrawn) A method of disinfecting mammalian tissue comprising applying a tissue antiseptic composition of claim 41 to mammalian tissue.

Page 17 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

- 59. (Withdrawn) A method of delivering a pharmaceutical agent to a mammal comprising applying a transdermal drug delivery composition of claim 50 to mammalian skin.
- 60. (Withdrawn) A method of delivering a pharmaceutical agent to a mammal comprising applying a transdermal drug delivery composition of claim 51 to mammalian skin.
- 61. (Withdrawn) A method of delivering a pharmaceutical agent to a mammal comprising applying a transdermal drug delivery composition of claim 52 to mammalian skin.
- 62. 65. (Canceled)
- 66. (Previously Presented) A water-in-oil emulsion comprising:

  a vinyl polymer comprising the reaction product of monomers comprising:
  isooctyl acrylate, stearyl methacrylate, and polyethylene oxide methacrylate;
  an oil phase; and

. 1

a water phase.

67. (Previously Presented) A moisturing composition comprising a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising: isooctyl acrylate, stearyl methacrylate, and polyethylene oxide methacrylate;

an oil phase; and

a water phase.

Page 18 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

68. (Currently Amended) A <u>mammalian</u> tissue antiseptic composition comprising an antimicrobial agent and a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising: isooctyl acrylate, stearyl methacrylate, and polyethylene oxide methacrylate;

an oil phase; and

a water phase; and

an antimicrobial.

69. (Previously Presented) A personal care composition comprising a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising: isooctyl acrylate, stearyl methacrylate, and polyethylene oxide methacrylate;

an oil phase; and

a water phase.

70. (Previously Presented) A transdermal drug delivery composition comprising a pharmaceutical agent and a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising: isooctyl acrylate, stearyl methacrylate, and polyethylene oxide methacrylate;

an oil phase; and

a water phase; and

a pharmaceutical agent.

71. (Withdrawn) A water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising:

(i) polyethylene oxide methacrylate and (ii) isooctyl acrylate, 2-ethylhexyl acrylate, or

Page 19 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

both;

an oil phase; and a water phase.

72. (Withdrawn) A moisturing composition comprising a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising:
(i) polyethylene oxide methacrylate and (ii) isooctyl acrylate, 2-ethylhexyl acrylate, or

both;

an oil phase; and

a water phase.

73. (Withdrawn – Previously Presented) A tissue antiseptic composition comprising an antimicrobial agent and a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising:

(i) polyethylene oxide methacrylate and (ii) isooctyl acrylate, 2-ethylhexyl acrylate, or both;

an oil phase; and

a water phase.

74. (Withdrawn) A personal care composition comprising a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising:

(i) polyethylene oxide methacrylate and (ii) isooctyl acrylate, 2-ethylhexyl acrylate, or both;

an oil phase; and

a water phase.

Page 20 of 24

Serial No.: 09/966,511 Confirmation No.: 4740 Filed: 28 September 2001

For: WATER-IN-OIL EMULSIONS WITH ETHYLENE OXIDE GROUPS, COMPOSITIONS, AND METHODS

75. (Withdrawn – Previously Presented) A transdermal drug delivery composition comprising a pharmaceutical agent and a water-in-oil emulsion comprising:

a vinyl polymer comprising the reaction product of monomers comprising:

(i) polyethylene oxide methacrylate and (ii) isooctyl acrylate, 2-ethylhexyl acrylate, or both;

an oil phase; and a water phase.